

STRESS DISTRIBUTION ANALYSIS OF 3D SOLID MODEL OF MAXILLA DUE TO BRUXISM USING FINITE ELEMENT METHOD: RECONSTRUCTING MODEL FROM CBCT RADIOGRAPH OF HUMAN MAXILLA

Anthony Sugiharta Budiawan*, Aldilla Miranda**, Andi Isra Mahyuddin*, Tatacipta Dirgantara*, Sandro Mihradi*, Sri Wendari**, Nunung Rusminah**

* Faculty of Mechanical and Aerospace Engineering, Institut Teknologi Bandung

** Periodontology Department, Faculty of Dentistry, Universitas Padjadjaran

ABSTRACT

Objectives: To construct three dimensional (3D) solid model of maxilla based on CBCT's radiograph and then to analyze the stress distribution on tooth and periodontal structure of maxilla due to bruxism using finite element method.

Material and Method: 3D solid model of maxilla in this study include the teeth, periodontal ligaments, cortical bones, and trabecular bones. The basic model was constructed from CBCT radiograph of human maxilla and improved to be a 3D solid model. To simulate bruxing force, the load were applied to the model with the initial force magnitudes of 570 N for the incisive-canine tooth and 911 N for the premolar-molar tooth as the loading condition.

Result: The principal stress from each movement with a maximum of 200 MPa on the jaw bones, 160 MPa on the teeth, and 80 MPa on the periodontal ligaments.

Conclusions: The highest stress appeared to be found at jaw bones, the result of this study will aid dentists who are attempting to identify the location and magnitude of stress concentration caused by bruxism. Afterwards this information can be used as a source of consideration in making splint or other protocol therapy, also it can be used as a reference for the next step of research in bruxism.

INTRODUCTION

About 45 million people in United States have a problem in parafunctional habit like bruxism.¹ Due to the parafunctional habit that occurs in long term are tooth attrition, widening the periodontal membrane, alveolar bone resorption, and even led to disruption of temporomandibular joint. It is caused by the presence of the load exceeds normal occlusion and continuously distributed from the surface of teeth to periodontal tissues.^{2,3,4} There has been research on the movement patterns of bruxism, but there has not been investigated regarding the stress distribution due to traumatic occlusion caused by bruxism.⁵ Therefore, researchers are interested in conducting research on stress distribution analysis of 3D solid model of maxilla due to bruxism using finite element method. Reconstruction of finite element model of human maxilla or mandible now becomes more accurate and simple with the CBCT radiograph.

RESULT AND DISCUSSION




Figure 6. Maximum stress distribution on maxilla for IC movement




Figure 7. Maximum stress distribution on maxilla for ICP movement





Figure 8. Maximum stress distribution on maxilla for ICPM movement



Figure 9. Maximum stress distribution on



Certificate

This is to certify that

Aldilla Miranda

**Participated in the 10th
Asian Congress
of Oral and Maxillo-Facial Radiology
As
Lecture**

**Inna Grand Bali Beach Hotel
Sanur, Bali - Indonesia
November 20 - 22, 2014**



SUHARDJO SITAM

CHAIRMAN OF THE 10TH ACOMFR 2014

Akreditasi PB PDGI
Accredited by Indonesian Dental Association
No:SKP-I/034rev/PB PDGI/VII/2014
Cakupan International
International Coverage

Peserta/Participant: 7,5 SKP
Pembicara Oral/Oral Presentation: 3 SKP
Pembicara Poster/Poster Presentation: 4 SKP
Juri Poster/Poster Jury: 5 SKP
Moderator/Moderator: 3 SKP
Panitia/Committee: 3 SKP